

Resolution of the $B \rightarrow \pi\pi$, πK puzzles

Hsiang-nan Li¹ and Satoshi Mishima²

¹*Institute of Physics, Academia Sinica, Taipei, Taiwan 115, Republic of China,*

¹*Department of Physics, Tsing-Hua University, Hsinchu, Taiwan 300, Republic of China,*

¹*Department of Physics, National Cheng-Kung University, Tainan, Taiwan 701, Republic of China and*

²*Theory Group, Deutsches Elektronen-Synchrotron DESY, 22607 Hamburg, Germany*

We show that there exist uncanceled soft divergences in the k_T factorization for nonfactorizable amplitudes of two-body nonleptonic B meson decays, similar to those identified in hadron hadroproduction. These divergences can be grouped into a soft factor using the eikonal approximation, which is then treated as an additional nonperturbative input in the perturbative QCD formalism. Viewing the special role of the pion as a $q\bar{q}$ bound state and as a pseudo Nambu-Goldstone boson, we postulate that the soft effect associated with it is significant. This soft factor enhances the nonfactorizable color-suppressed tree amplitudes, such that the branching ratios $B(\pi^0\pi^0)$ and $B(\pi^0\rho^0)$ are increased under the constraint of the $B(\rho^0\rho^0)$ data, the difference between the direct CP asymmetries $A_{CP}(\pi^\mp K^\pm)$ and $A_{CP}(\pi^0 K^\pm)$ is enlarged, and the mixing-induced CP asymmetry $S_{\pi^0 K_S}$ is reduced. Namely, the known $\pi\pi$ and πK puzzles can be resolved simultaneously.

PACS numbers: 13.25.Hw, 12.38.Bx, 12.39.St

The more precise data of the $B \rightarrow \pi\pi$, πK decays have sharpened the discrepancies with the theoretical predictions from the factorization approaches, such as the perturbative QCD (PQCD) approach based on the k_T factorization theorem [1, 2]. The observed $B^0 \rightarrow \pi^0\pi^0$ branching ratio [3] remains several times larger than the naive expectation. The direct CP asymmetry of the $B^\pm \rightarrow \pi^0 K^\pm$ decays differs dramatically from that of $B^0 \rightarrow \pi^\mp K^\pm$ decays. There is a deviation between the extractions of the standard model parameter $\sin(2\phi_1)$ from the penguin-dominated $B^0 \rightarrow \pi^0 K_S$ modes and from the tree-dominated $b \rightarrow c\bar{c}s$ modes. All these discrepancies are closely related to the color-suppressed tree amplitudes C [4]. The $B^0 \rightarrow \pi^0\rho^0$ branching ratios from PQCD and QCD factorization (QCDF), being sensitive to C , are also much lower than the data [5, 6]. However, the estimate of C from PQCD is well consistent with the measured $B^0 \rightarrow \rho^0\rho^0$ branching ratio [7]. Proposals resorting to new physics [8] mainly resolve the πK puzzle without addressing the peculiar feature of C in the $\pi^0\pi^0$, $\pi^0\rho^0$, and $\rho^0\rho^0$ modes, while those to QCD effects are usually strongly constrained by the $\rho\rho$ data [9]. It indicates the difficulty of resolving the $\pi\pi$ and πK puzzles simultaneously.

The color-suppressed tree amplitude C seems to be an important but the least understood quantity in B meson decays. Viewing that all the puzzles appear in the C -sensitive quantities, we shall carefully investigate QCD effects on C , and their impact on the $B \rightarrow \pi\pi$, πK decays. Once a mechanism identified for C respects the conventional factorization theorem, it is unlikely to be a resolution due to the $B \rightarrow \rho\rho$ constraint mentioned above [7]. That is the reason the higher-order corrections calculated in QCDF [10],




FIG. 1: LO diagrams for a nonfactorizable amplitude.

which obey the collinear factorization, cannot resolve the $\pi\pi$ puzzle. It has been pointed out by Collins and Qiu [11] that the k_T factorization breaks down in complicated QCD processes like high- p_T hadron hadroproduction. To factorize the collinear gluons associated with, say, one of the initial-state hadrons, one needs to eikonalize the valence quark lines to which the collinear gluons attach. Those eikonal lines, i.e., Wilson lines from another initial-state hadron and the final-state hadrons, should cancel in order to have the universality of the considered parton distribution function. However, the required cancellation is not exact in the k_T factorization, though it is in the collinear factorization. The k_T factorization still holds for simple processes like deeply inelastic scattering (DIS), which does not involve the Wilson lines from the other hadrons.

The above observation provides a clue for resolving the $\pi\pi$ and πK puzzles. It is easy to see that a factorizable amplitude, involving only a B meson transition form factor, mimics simple DIS, and a nonfactorizable[27] amplitude, involving dynamics of three hadrons, mimics the complicated hadron




FIG. 2: NLO diagrams for Fig. 1(a) that are relevant to the factorization of the M_2 meson wave function.

hadroproduction. The k_T factorization for a factorizable B meson decay amplitude has been proved [12]. The k_T factorization for a nonfactorizable amplitude has not, though it has been widely employed in the PQCD analysis. Below we shall identify the residual infrared divergence in the k_T factorization for a nonfactorizable amplitude at one loop. Contrary to high- p_T hadron hadroproduction, this residual infrared divergence can be factorized into a soft factor in two-body nonleptonic B meson decays, following the procedure in [13], so that the universality of a k_T -dependent meson wave function is restored. A nonfactorizable amplitude then remains calculable in the PQCD approach after parameterizing the soft factor.

Consider the $B(P_B) \rightarrow M_1(P_1)M_2(P_2)$ decay, where $P_{B,1,2}$ represent the momenta of the B , M_1 , and M_2 mesons, respectively. Start with the leading-order (LO) nonfactorizable emission diagram in Fig. 1(a) resulting from the operator O_2 [14]. We add a radiative gluon of momentum l emitted by the valence quark in M_2 . The attachment of the radiative gluon to the b quark line shown in Fig. 2(a) leads to a Wilson line from infinity to the origin, i.e., the weak vertex.




FIG. 3: Two-particle reduced NLO diagrams for the M_2 meson wave function.

This piece is factorized in color flow by itself with the color factor C_F . The attachment to the hard gluon in Fig. 2(b) generates two Wilson lines, one of which runs from the position z_2 of the valence anti-quark in M_2 to infinity. The attachments to the virtual anti-quark in Fig. 2(c) and to the valence quark in the M_1 meson in Fig. 2(d) also generate the Wilson line running from z_2 to infinity. The combination of these three pieces with the same Wilson line is factorized in color flow. As to the next-to-leading-order (NLO) two-particle reducible diagrams, such as the self-energy correction to the valence quark in Fig. 3(a) and the gluon exchange between the valence quark and the valence anti-quark in Fig. 3(b), their factorization into the M_2 meson wave function is straightforward.

The other attachments, such as Figs. 2(e) and 2(f), should cancel in order to have the universality of the M_2 meson wave function. Eikonalizing the spectator quark line on the B meson side, Fig. 2(e) gives the integral

$$I_E = C_F \int \frac{d^4 l}{(2\pi)^4} \text{tr} \left[\dots \frac{-i(\not{P}_2 - \not{k}_2 + \not{k}_1 - \not{l})}{(P_2 - k_2 + k_1 - l)^2 + i\epsilon} \right. \\ \times (-ig\gamma_\beta)\gamma_5 \not{P}_2 (-ig\gamma_\alpha) \frac{i(\not{k}_2 + \not{l})}{(k_2 + l)^2 + i\epsilon} \left. \right] \\ \times \frac{-gn_+^\alpha}{n_+ \cdot l + i\epsilon} \frac{-i}{l^2 + i\epsilon}, \quad (1)$$

where the vector $n_+ = (1, 0, 0_T)$ is along the momentum P_1 , the $\dots$ denotes the rest of the integrand, and $\gamma_5 \not{P}_2$ comes from the twist-2 structure of the M_2 meson wave function. Another piece from Fig. 2(b) contains the Wilson line running from the position of the spectator in the M_1 meson to infinity. This piece, together with Fig. 2(f), leads to the loop integral the same as Eq. (1), but with the eikonal propagator being replaced by $1/(-n_+ \cdot l + i\epsilon)$. Employing the principal-value prescription

$$\frac{1}{n_+ \cdot l + i\epsilon} + \frac{1}{-n_+ \cdot l + i\epsilon} = -2\pi i \delta(l^-), \quad (2)$$

we find a NLO residual soft divergence from the

Glauber region with $l^- = 0$, which seems to violate the universality of the M_2 meson wave function.

The residual soft divergence may not cause trouble, if the LO amplitude $\mathcal{M}^{(0)}$ is real. Expanding the decay width up to NLO, we have

$$|\mathcal{M}|^2 = |\mathcal{M}^{(0)}|^2 + 2\text{Re}[\mathcal{M}^{(0)}\mathcal{M}^{(1)*}]. \quad (3)$$

According to Eq. (8), the residual soft divergence will be purely imaginary, if $\mathcal{M}^{(0)}$ is real, so it does not survive in the second term $\text{Re}[\mathcal{M}^{(0)}\mathcal{M}^{(1)*}]$. This is the reason the diagrams Fig. 2 are calculable in the collinear factorization, and result is mainly real in QCDF [10]. $\mathcal{M}^{(0)}$ is complex in the k_T factorization, because the partons carry transverse momenta, and the internal lines can go on mass shell at finite momentum fractions [15]. Then the residual soft divergence contributes to $\text{Re}[\mathcal{M}^{(0)}\mathcal{M}^{(1)*}]$. It has been known that the end-point singularity from a vanishing momentum fraction is developed at twist 3 in QCDF for a nonfactorizable decay amplitude. The presence of the end-point singularity signals the inappropriate theoretical framework, while the residual soft divergence in the k_T factorization for a nonfactorizable B meson decay amplitude can be factorized and be handled by introducing an additional nonperturbative input as shown below.

We eikonalize the valence quark line of the M_2 meson according to the observation in [13]: for P_2 in the minus direction, the poles

$$l^+ = \frac{|\mathbf{k}_{2T} + \mathbf{l}_T|^2}{2k_2^-} - i\epsilon, \quad (4)$$

$$l^+ = k_1^+ - \frac{|\mathbf{k}_{1T} - \mathbf{k}_{2T} - \mathbf{l}_T|^2}{2(P_2^- - k_2^-)} + i\epsilon, \quad (5)$$

from the denominators $(k_2 + l)^2 + i\epsilon$ and $(P_2 - k_2 + k_1 - l)^2 + i\epsilon$ in Eq. (1), respectively, are apart from each other by $O(k_1^+)$. As long as k_1^+ is greater than $O(\Lambda_{\text{QCD}})$, Λ_{QCD} being the QCD scale, we can deform the contour of l^+ , such that l^+ remains larger than $O(\Lambda_{\text{QCD}})$, and the approximation

$$k_2^- l^+ > O(m_B \Lambda_{\text{QCD}}) \gg |\mathbf{k}_{2T} + \mathbf{l}_T|^2 \sim O(\Lambda_{\text{QCD}}^2), \quad (6)$$

holds. We then have the eikonalization of the valence quark line even in the Glauber region,

$$\dots \gamma_5 \not{P}_2 \not{k}_2 + \not{l} \frac{1}{(k_2 + l)^2 + i\epsilon} \approx \dots \gamma_5 \not{P}_2 \frac{1}{l^+ + i\epsilon}. \quad (7)$$

Picking up the pole $l^+ = 0 - i\epsilon$ from the above eikonal propagator, the residual soft divergence in Fig. 2 is factorized into the convolution

$$I_a^{(1)} \approx i \frac{\alpha_s}{\pi} C_F \int \frac{d^2 l_T}{l_T^2} \mathcal{M}_a^{(0)}(\mathbf{l}_T), \quad (8)$$




FIG. 4: A NLO diagram that does not contribute the residual soft divergence.

where $\mathcal{M}_a^{(0)}$ denotes the LO amplitude from Fig. 1(a).

If the radiative gluon is emitted by the valence anti-quark of the M_2 meson in Fig. 1(a), the corresponding loop integral does not generate the residual soft divergence. Take the NLO diagram in Fig. 4 as an example, where the radiative gluon attaches to the spectator of the M_1 meson. All the poles of l^+ are located in the lower half plane, and not pinched, so the integral diminishes. That is, these diagrams contribute only the infrared divergence that ought to be absorbed into the M_2 meson wave function.

Adding the second radiative gluon emitted by the valence quark in the M_2 meson, we have

$$I_a^{(2)} \approx \frac{1}{2} \left(i \frac{\alpha_s}{\pi} C_F \right)^2 \int \frac{d^2 l_{1T} d^2 l_{2T}}{l_{1T}^2 l_{2T}^2} \mathcal{M}_a^{(0)}(\mathbf{l}_{1T} + \mathbf{l}_{2T}), \quad (9)$$

whose derivation is similar to the two-loop analysis in [11]. Extending the analysis to higher orders, and neglecting the dependence of $\mathcal{M}_a^{(0)}$ on the transverse loop momenta $\mathbf{l}_T$, the divergence can be summed into an exponential factor, $I_a \approx e^{iS_e} \mathcal{M}_a^{(0)}$ with S_e being a parameter. Considering the soft corrections to the LO diagram in Fig. 1(b) from the gluons radiated by the valence anti-quark in M_2 , we obtain $I_b \approx e^{-iS_e} \mathcal{M}_b^{(0)}$, where the minus sign is attributed to the radiation from the anti-quark. The above modified k_T factorization formalism also applies to nonfactorizable amplitudes from other tree and penguin operators, and to nonfactorizable annihilation amplitudes.

The soft factor has a dynamical origin similar to that of a meson wave function: the former (latter) absorbs the soft (collinear) gluons. The behavior of the soft factor can be obtained either by nonperturbative methods or from experimental data. We then study its impact on nonleptonic two-body B meson decays. The color-suppressed tree amplitudes are small at LO due to the small Wilson coefficient a_2 for the factorizable contribution and to the pair cancellation between Figs. 1(a) and 1(b) for the nonfactorizable contribution. The presence of the soft factor can convert the destructive interference in Fig. 1 into a constructive one, leading to strong enhancement. Its effect on

the color-allowed tree and penguin (including annihilation) amplitudes should be minor, which are dominated by factorizable contributions.

We seek experimental constraints on the soft factor by comparing the data [3] (in units of 10^{-6})

$$\begin{aligned} B(\pi^0\pi^0) &= 1.55 \pm 0.19, [(0.29^{+0.50}_{-0.20})] \\ B(\pi^0\rho^0) &= 2.0 \pm 0.5, [\approx 0.7] \\ B(\rho^0\rho^0) &= 0.74^{+0.30}_{-0.27}, [(0.92^{+1.10}_{-0.56})], \end{aligned} \quad (10)$$

with the NLO PQCD predictions in the square brackets, which are quoted from [16], [17], and [7], respectively. The results from QCDF [6, 18] are similar. The above comparison motivates us to postulate that the soft effect is significant (negligible) in the decays with M_2 being a pseudoscalar (vector) meson. That is, we associate a soft factor with $M_2 = \pi, K$, but not with $M_2 = \rho$ (the soft effect associated with the kaon is not crucial actually). Because the $B^0 \rightarrow \pi^0\rho^0$ decay involves both types of amplitudes with the pion and the ρ meson as M_2 , it is natural that the discrepancy is in between as indicated by Eq. (10). The larger soft effect from the multi-parton states in the pion than in the ρ meson can be understood by means of the simultaneous role of the pion as a $q\bar{q}$ bound state and as a Nambu-Goldstone (NG) boson [19]: the valence quark and anti-quark of the pion are separated by a short distance, like those of the ρ meson, in order to reduce the confinement potential energy. The multi-parton states of the pion spread over a huge space-time in order to meet the role of a massless NG boson.

The factorization formulas for the $B \rightarrow \pi\pi, \pi\rho, \pi K$, and ρK decays can be found in [16, 20]. According to our derivation, we multiply the b quark nonfactorizable emission amplitudes, both tree and penguin, by e^{iS_e} (e^{-iS_e}) with the hard gluon being emitted by the valence anti-quark (quark) in M_2 . The dependence on S_e of those C -sensitive quantities is displayed in Fig. 5. The branching ratios $B(\pi^0\pi^0)$ and $B(\pi^0\rho^0)$ grow quickly with decreasing S_e from the NLO PQCD values in Eq. (10), and become close to the data when S_e reaches $-\pi/2$. The direct CP asymmetry $A_{CP}(\pi^0 K^\pm)$ increases from the NLO PQCD result around -0.01 [16] to above 0.05 for $S_e < -\pi/4$, whose agreement with the data $A_{CP}(\pi^0 K^\pm) = 0.050 \pm 0.025$ [3] is satisfactory. The deviation of the mixing induced CP asymmetry $\Delta S_{\pi^0 K_S} \equiv S_{\pi^0 K_S} - S_{c\bar{c}s}$ descends from the NLO PQCD value $+0.07$ to -0.04 for $S_e = -\pi/2$. Compared to the data $S_{\pi^0 K_S} = 0.57 \pm 0.17$ and $S_{c\bar{c}s} = 0.672 \pm 0.024$ [3], the consistency has been improved. The ratio $C/T = 0.53e^{-2.2i}$ for the $B \rightarrow \pi\pi$ decays, corresponding to $S_e \approx -\pi/2$, is consistent with the extraction in [4], T being the color-allowed tree amplitude. An equivalent viewpoint is that the

$B(\pi^0\pi^0)$ data constrain $S_e \sim -\pi/2$, which then leads to the predictions for other quantities in Fig. 5.

We have confirmed that $B(\pi^\mp\pi^\pm)$, $B(\pi^0\pi^\pm)$ and all $B(\pi K)$ change slightly from those in [16], since they are less sensitive to C . $A_{CP}(\pi^\mp K^\pm)$ remains around -0.1 [1] for arbitrary S_e , and in agreement with the data $A_{CP}(\pi^\mp K^\pm) = -0.098^{+0.012}_{-0.011}$ [3]. The small variation of the curve is attributed to the soft effect on the nonfactorizable color-allowed and penguin contributions. $S_{\rho^0 K_S}$ does not change much, because $M_2 = \rho$ in this case, and the involved C is not modified. The NLO PQCD prediction $\Delta S_{\rho^0 K_S} \approx -0.15$ [20] is consistent with the data $S_{\rho^0 K_S} = 0.63^{+0.17}_{-0.21}$ [3]. For those penguin-dominated two-body modes without involving C , like $B \rightarrow \phi K$, their mixing-induced CP asymmetries are not affected either.

Our resolution differs from those based on new physics models, such as the fourth-generation model [21, 22], in which it is the electroweak penguin amplitude that is enhanced. These proposals, in which new weak phases are introduced, change $S_{\rho^0 K_S}$ and $S_{\phi K_S}$. Our proposal, which does not involve multiple intermediate states, differs from the elastic rescattering models for final-state interaction [23]. A large C has been generated through the charge exchange mechanism in [24]. Since the QCDF approach was employed, the parameter scenario ‘‘S4’’ [6] or the inelastic scattering [25] has to be incorporated in order to get the correct result for $A_{CP}(\pi^\mp K^\pm)$. The exchange mechanism, which requires turning of two energetic quarks into opposite directions, is suppressed according to the factorization theorem. Moreover, $\Delta S_{\pi^0 K_S}$ remains positive in [24].

In this letter we have identified the uncanceled soft divergences in the k_T factorization for the nonfactorizable B meson decay amplitudes, which are similar to those existent in hadron hadroproduction. The divergences are factorizable and demand the introduction of the additional soft factor, under which we have computed C in the PQCD approach, and found a simultaneous resolution of all the puzzles: $B(\pi^0\pi^0)$ and $B(\pi^0\rho^0)$ are enhanced, the difference between $A_{CP}(\pi^\mp K^\pm)$ and $A_{CP}(\pi^0 K^\pm)$ is enlarged, and $\Delta S_{\pi^0 K_S}$ is reduced for the single soft parameter S_e around $-\pi/2$. The constraint on C from the $B \rightarrow \rho\rho$ data is evaded, because of the special role of the pion as a $q\bar{q}$ bound state and as a pseudo NG boson. The mechanism identified here can be verified or falsified by more precise data in the future. Based on our observation, we agree that the $B \rightarrow \pi K$ data have not yet revealed a new physics signal [26].

This work was supported by the National Center for Theoretical Sciences and National Science Council of R.O.C. under Grant No. NSC-95-2112-M-050-MY3.




FIG. 5: S_e dependence of (a) $B(\pi^0\pi^0)$ and $B(\pi^0\rho^0)$ in units of 10^{-6} , of (b) $A_{CP}(\pi^0K^\pm)$ and $A_{CP}(\pi^\mp K^\pm)$, and of (c) $\Delta S_{\pi^0K_S}$ and $\Delta S_{\rho^0K_S}$. The data (horizontal bands) for ΔS are not shown due to their large errors.

-
- [1] Y.Y. Keum, H-n. Li, and A.I. Sanda, Phys. Lett. B **504**, 6 (2001); Phys. Rev. D **63**, 054008 (2001).
- [2] C.D. Lü, K. Ukai, and M.Z. Yang, Phys. Rev. D **63**, 074009 (2001).
- [3] Heavy Flavor Averaging Group, E. Barberio et al., arXiv:0808.1297 [hep-ex].
- [4] C. Chiang et al., Phys. Rev. D **70**, 034020 (2004); Y.Y. Charng and H-n. Li, Phys. Rev. D **71**, 014036 (2005); R. Fleischer, S. Recksiegel, F. Schwab, Eur. Phys. J. C **51**, 55 (2007).
- [5] C.D. Lü and M.Z. Yang, Eur. Phys. J. C **23**, 275 (2002).
- [6] M. Beneke and M. Neubert, Nucl. Phys. **B675**, 333 (2003).
- [7] H-n. Li and S. Mishima, Phys. Rev. D **73**, 114014 (2006).
- [8] R. Fleischer, S. Jager, D. Pirjol, and J. Zupan, arXiv:0806.2900 [hep-ph], and references therein.
- [9] M. Beneke, J. Rohrer, and D. Yang, Nucl. Phys. **B774**, 64 (2007).
- [10] M. Beneke and D. Yang, Nucl. Phys. **B736**, 34 (2006); M. Beneke and S. Jager, Nucl. Phys. **B751**, 160 (2006); G. Bell, Nucl. Phys. **B795**, 1 (2008); V. Pilipp, Nucl. Phys. **B794**, 154 (2008).
- [11] J. Collins and J.W. Qiu, Phys. Rev. D **75**, 114014 (2007); J. Collins, arXiv:0708.4410 [hep-ph].
- [12] M. Nagashima and H-n. Li, Phys. Rev. D **67**, 034001 (2003).
- [13] C.P. Chang and H-n. Li, arXiv:0904.4150 [hep-ph].
- [14] G. Buchalla, A. J. Buras, and M. E. Lautenbacher, Review of Modern Physics, **68**, 1125 (1996).
- [15] J. Chay, H-n. Li, and S. Mishima, Phys. Rev. D. **78**, 034037 (2008).
- [16] H-n. Li, S. Mishima, and A.I. Sanda, Phys. Rev. D **72**, 114005 (2005).
- [17] H-n. Li and S. Mishima, in preparation.
- [18] M. Beneke, J. Rohrer and D. Yang, Nucl. Phys. **B774**, 64 (2007); H.Y. Cheng and K.C. Yang, Phys. Rev. D **78**, 094001 (2008).
- [19] G.P. Lepage and S.J. Brodsky, Phys. Lett. B **87**, 359 (1979); S. Nussinov and R. Shrock, arXiv:0811.3404 [hep-ph]; M. Duraisamy and A.L. Kagan, arXiv:0812.3162 [hep-ph].
- [20] H-n. Li and S. Mishima, Phys. Rev. D **73**, 094020 (2006).
- [21] W.S. Hou, H-n. Li, S. Mishima, and M. Nagashima, Phys. Rev. Lett. **98**, 131801 (2007).
- [22] A. Soni et al., arXiv:0807.1971 [hep-ph].
- [23] C.K. Chua, W.S. Hou, and K.C. Yang, Phys. Rev. D **65**, 096007 (2002); A.B. Kaidalov and M.I. Vysotsky, Phys. Lett. B **652**, 203 (2007); M.I. Vysotsky, arXiv:0901.2245; A.F. Falk et al., Phys. Rev. D **57**, 4290 (1998).
- [24] C.K. Chua, Phys. Rev. D **78**, 076002 (2008).
- [25] H.Y. Cheng, C.K. Chua, and A. Soni, Phys. Rev. D **71**, 014030 (2005).
- [26] M. Ciuchini et al., arXiv:0811.0341; N. Mahajan, arXiv:0812.0230 [hep-ph].
- [27] Here a "nonfactorizable" amplitude refers to a contribution that does not respect the naive factorization assumption, which can also be called as a "spectator" amplitude.